



<https://www.biodiversitylibrary.org/>

Proceedings of the Zoological Society of London

London, Academic Press, [etc.], 1833-1965

<https://www.biodiversitylibrary.org/bibliography/44963>

pt.12-15 (1844-1847): <https://www.biodiversitylibrary.org/item/46217>

Article/Chapter Title: Sowerby 1846

Page(s): Page 91, Page 92, Page 93, Page 94, Page 95, Page 96, Page 97

Holding Institution: Natural History Museum Library, London

Sponsored by: Natural History Museum Library, London

Generated 16 July 2023 9:12 PM

<https://www.biodiversitylibrary.org/pdf4/1603482i00046217.pdf>

This page intentionally left blank.

"Descriptions of thirteen new species of *Brachiopoda.*" By G. B. Sowerby, F.L.S.

TEREBRATULA NIGRICANS. *Ter. testā anticē rotundatā, posticē acuminatā, tenuiusculā, nigricante; valvis inæqualibus, radiatim costatis, costis rotundatis; lined marginali rectiusculā; valvā dorsali subtrigond, depressiusculā, rotundatā, lateribus posticis declivibus; foramine magno, haud integro, quadrato; ared cardinali magnā, planiusculā, lateribus rotundatis; deltidiis angustis ad latera foraminis coalescentibus; valvā ventrali depressā, transversim ovatā; dentibus cardinalibus ut in *T. psittacea*; margine valvarum crenulato.*

A single specimen of this very interesting species was found in the collection of the late G. Humphrey, without locality. This and the *T. psittacea* are the only species that are not punctated.

TEREBRATULA JAPONICA. *Ter. testā oblongā, anticē rotundatā, tenui, albicante; valvis subæqualiter convexis, longitudinaliter radiatim striatis, striis numerosis, subirregularibus, subbifurcatis, ad latera prope cardinem confertiusculis; lateribus prope cardinem subplanulatis; lined marginali rectiusculā, ad latera posticē declivi; valvā dorsali posticē subproductā, truncatā; foramine mediocri, haud integro, perobliquo; ared cardinali inconspicuā, deltidiis obsoletis; interno parvo, $\frac{1}{3}$ longitudinis valvæ, ramulos duos angustos, demum amentum latum flexuosum sistente; marginibus valvarum minutissimè denticulato.*

Shell oblong, rounded in front, thin, whitish; valves nearly equally convex, longitudinally radiately striated, striae numerous, rather irregularly dichotomous and very close-set on the sides near the hinge; the sides near the hinge rather flattened; marginal line nearly straight, inclining to the sides near the hinge; dorsal valve somewhat produced posteriorly and truncated, with a moderate-sized, very oblique and incomplete perforation; cardinal area indistinct, with obsolete deltidia; internal appendages small, one-third the length of the valve, with two narrow little branches, and then a broad flexuous loop; margin of the valves very minutely denticulated.

In Mr. Cuming's collection, from Japan. Easily distinguished from *T. cancellata* of Koch by its foramen not being entire.

TEREBRATULA CRENULATA. *Ter. testā suborbiculari, posticē subacuminatā, subtruncatā, crassiusculā, albicante; valvis inæqualibus, radiatim costatis, costis paucis, majusculis, rotundatis; lined marginali flexuosa, anticē subsinuatā; foramine magno, subintegro; ared cardinali magnā, subplanulatā, margine undulato; deltidiis magnis, discretis; valvæ ventralis margine postico ad utrumque latus declivi; ossiculo interno e spindā validā obtusā anticē porrectā, ramulis duobus lateralibus retroversis; margine valvarum crenato.*

Shell suborbicular, rather attenuated and subtruncate behind, rather thick and whitish; valves unequal, with few rather large, rounded, radiating ribs; marginal line flexuous, slightly sinuated in

front; perforation large, nearly entire; cardinal area large, flattish, with an undulated margin; deltidia large, separate; posterior margin of the ventral valve inclined on either side; internal appendage consisting of a single strong spine standing forward, and with two lateral reflected branches; margin of the valves crenated.

From Santa Cruz, in Mr. Cuming's collection.

TEREBRATULA ROSEA, Humphrey. *Ter. testa oblongo-ovali, subdepressa, anticè subattenuata, crassata, rosata, albicante radiatim picta; valvis subæqualibus, lœvibus; linea marginali subflexuosa, anticè reflexa; valva dorsali majori, posticè subproducta, truncata; foramine minimo, integro; area cardinali latiuscula, deltidis coalescentibus, sulco mediano discretis; valva ventrali ovali, anticè paululum rotundato-attenuata; linea marginali ex umbone ad utrumque latus declivi; ossiculo costata elevata, porrecta, simplici constante; margine valvarum integro.*

Shell of an oblong-oval form, rather depressed, slightly attenuated behind, thick, rose-red, painted with paler radiating marks; valves nearly equal, smooth; marginal line somewhat flexuous, reflected in front; dorsal valve the larger, rather produced posteriorly and truncated; perforation very small, entire; cardinal area rather wide, with united deltidia marked by a mesial groove; ventral valve oval, anteriorly slightly attenuated and rounded, its marginal line slanting downwards from the apex on each side; internal appendage consisting of a single elevated rib standing out; margin of the valves entire.

From Brazil, according to the late G. Humphrey. In Mr. Cuming's and other collections.

TEREBRATULA RUBICUNDA. (*T. sanguinea*, Quoy, Astr.) *Ter. testa suborbiculari, posticè subacuminata, gibba, glabra, rubra, valvis inæqualibus, subirregularibus; linea marginali ad latera subflexuosa, anticè sinuata; valva dorsali posticè producta, truncata; foramine magno, obliqua, subintegro; area cardinali rotundata; deltidis magnis, distinctis; carina dorsali latâ, prominente, utrinque rotundato-angulata, obtusa; valva ventrali subpentagonalis, posticè angustiori, medianè latâ, rotundato-subangulata; anticè subtruncata, sulco mediano lato, conspicuo; ossiculo interno magno, ferè ut in *T. dorsata* efformato; margine valvarum integro.*

Shell nearly orbicular, rather acuminate posteriorly, gibbous, smooth, of a red colour; valves unequal, rather irregular; marginal line slightly flexuous on the sides, sinuated in front; dorsal valve produced behind, truncated, with a large, oblique, nearly entire perforation; cardinal area rounded, deltidia large, separate; mesial ridge broad, prominent, angularly rounded, and obtuse on both sides; ventral valve somewhat pentagonal, narrow posteriorly, broad in the middle, with rounded angles, and slightly truncated in front; mesial groove broad and distinct; internal appendage as in *T. dorsata*; margin of the valves entire.

From the Moluccas; in Mr. Cuming's collection and in the British Museum.

TEREBRATULA SANGUINEA (*sanguinea*, Chemn.; *T. erythroleuca* of Quoy). *Ter. testa suborbiculari, anticè subemarginatā, gibbosiusculd, tenui, sanguined, radiis maculisque radiantibus albidis ornatā; linea marginali recta, anticè subsinuatā; valva dorsali posticè subacuminatā, truncatā, foramine mediocri, integro, areā cardinali latiusculd, marginibus subacutis, deltidis majusculis, coalescentibus; valva ventrali depresso-depressiusculd, transversim obovata, anticè subsinuatā, ossiculo interno primum format radios duos, deinde annulum centralem et amenta duo lateralia, demūm amentum superum integrum.*

Shell suborbicular, slightly notched in front, rather gibbous, thin, of a bright light red colour, with white rays and radiating spots; marginal line straight, slightly sinuated in front; dorsal valve rather acuminated behind and truncated; perforation middle-sized, complete; cardinal area rather broad, with sharpish edges; deltidia rather large and united; ventral valve somewhat depressed, transversely obovate, slightly sinuated in front; the internal appendage at first forms two rays, then a central ring and two lateral loops, and at length a reflected dorsal loop united to the central ring; margin of the valves entire.

From the island of Zebu, attached to coral under stones; H. Cuming.

TEREBRATULA INCONSPICUA. *Ter. testa rotundato-subtrigonali, posticè acuminato-rotundatā, anticè subsinuatā, obscurè rufa; valvis inæqualibus, glabris; linea marginali flexuosa; valva dorsali rotundato-subtrigona, maximā incompletā; areā cardinali latā, ad utrumque latus declivi; deltidis mediocribus, latè discretis; valva ventrali transversim oblonga, subplanulata, sulco mediano, lato, subinconspicuo; margine valvarum integro.*

Shell rounded, subtrigonal, acuminated and rounded behind, slightly sinuated before, dull red; valves unequal, smooth; marginal line flexuous; perforation large, incomplete; cardinal area broad, inclining on each side; deltidia of moderate size, widely separated; ventral valve transversely oblong, somewhat flattened, with a broad, rather indistinct mesial groove; margin of the valves entire.

From the late G. Humphrey's collection: locality unknown.

TEREBRATULA PULCHELLA. *Ter. testa subovata, posticè acuminato-rotundatā, laevi, albida, lineis nonnullis radiantibus rufis; valvis inæqualibus; linea marginali subflexuosa; valva dorsali subplanulata, posticè acuminata, anticè rotundata, foramine magno, incompleto; areā marginali inconspicua, lateribus rotundatis; deltidis parvis, discretis; valva ventrali subcirculari, planulata; ossiculo interno e gnomone porrecto, anticè posito, constante; margine valvarum integro.*

Shell subovate, acuminated and rounded behind, smooth, whitish with a few radiating red lines; valves unequal, marginal line somewhat flexuous; dorsal valve rather flattened, acuminated posteriorly, rounded in front; perforation large, incomplete; cardinal area indistinct, its sides rounded; deltidia small, separate; ventral valve

somewhat circular, flattened; internal appendage consisting of a single prominent gnomon near the front; margin of the valves entire.

Found by Mr. Cuming attached to corals at Calapan, isle of Mindoro; also from the island of Cocos, Lieut. Swainson; in the late G. Humphrey's collection.

TEREBRATULA COGNATA, Chemn. *Ter. testā subtrapezoidalē, anticē rotundatā, pallescente, nonnunquam rubente; valvis inæqualibus, radiatim obsoletē striatis, versus marginem oblitteratis; linea marginali lateraliter anticēque flexuosa; valvā dorsali convexā, līnd mediād inconspicuā; apice subacuminato reflexo; foramine magno, haud integro; areā cardinali angustā, deltidiis parvis, trigonalibus; valvā ventrali planata, margine postico rectiusculo; sulco mediano subperspicuo; ossiculo interno ramulos duos centrales, divergentes, ad apices expansos sistente; margine interno valvarum denticulato.*

Shell nearly trapezoidal, rounded in front, of a pale colour, sometimes reddish; valves unequal, obsoletely radiately striated, the striæ entirely obliterated near the margin; marginal line flexuous in front and on the sides; dorsal valve convex, with an inconspicuous central ridge, its apex somewhat acuminated, reflected, with a large incomplete perforation; cardinal area narrow, with small triangular deltidia; ventral valve flattened, its posterior margin nearly straight, with a scarcely evident central furrow; internal appendages consisting of two little central diverging branches, expanded at their apices; margin of the valves denticulated within.

There are two varieties in colour, from South Africa, according to the late G. Humphrey.

TEREBRATULA TRANSVERSA. *Ter. testā transversim subovatā, tenui, rudi, glabra, pallescente; linea marginali subflexuosa, anticē subsinuata; foramine maximo, incompleto; areā cardinali magnā, planata; deltidiis parvis, longè discretis; carina dorsali inconspicua, rotundata; valvā ventrali transversim oblongā, anticē rotundata, posticē in angulo obtusissimo desinente; sulco mediano subinconspicuo, rotundato; margine valvarum integro.*

Shell transversely subovate, thin, rugose, smooth, of a pale colour; marginal line somewhat flexuous, slightly sinuated in front; dorsal valve of a somewhat tetragonal ovate form, very obtusely angular behind and reflected; perforation very large, incomplete; cardinal area large and flattened; deltidia small, very distant; mesial ridge rounded, indistinct; ventral valve transversely oblong, rounded in front, and finishing in a very obtuse angle behind; mesial groove rounded; indistinct margin of the valves entire.

In Mr. Norris's collection and in that of Mr. Janelle.

TEREBRATULA RUBELLA, Sow. *Ter. testā subovatā, posticē subacuminata, subgibba, anticē subsinuata, rubra; valvis inæqualibus, glabris; linea marginali rectiuscula, anticē subsinuata; valvā dorsali posticē subacuminata, reflexa, carina medianā nullā, sulco mediano obsoletissimo, foramine parvo; areā cardinali angusta, ad*

latera rotundata, deltidiis majusculis, coalescentibus; valva ventrali ovata, sulco mediano latiusculo, emarginationem anticam efformante; ossiculo interno ut in T. australi; margine valvarum laevissimo.

Shell nearly oval, rather acuminated posteriorly, a little gibbous and slightly sinuated anteriorly; valves unequal, smooth; marginal line nearly straight, a little sinuated in front; dorsal valve rather acuminated posteriorly, reflected, without any mesial ridge, but with a very obsolete mesial furrow; perforation small; cardinal area narrow, rounded at the sides, with rather large united deltidia; ventral valve ovate, with a broad mesial groove forming a sinus in front; internal appendage as in *T. australis*.

From Japan. In Mr. Norris's and Mr. Cuming's collections.

TEREBRATULA LABRAPORENSIS. *Ter. testa suborbiculari, posticè acuminata, obtusa, crassiuscula, albida; valvis valde inæqualibus, radiatim obsoletè costatis; linea marginali subflexuosa; valva dorsali anticè rotundata, posticè acuminata, obtusa; foramine magno, integro; area cardinali magna, subplanulata; deltidiis majusculis, coalescentibus; carina dorsali inconspicua; valva ventrali suborbiculari, posticè subacuminata; margine valvarum crenulato.*

Shell suborbicular, acuminated behind, obtuse, thickish, whitish; valves very unequal, obsoletely radiately ribbed; marginal line somewhat flexuous; dorsal valve rounded in front, acuminated and obtuse behind; perforation large, entire; cardinal area large, somewhat flattened; deltidia rather large, united; mesial ridge indistinct; ventral valve nearly orbicular, slightly acuminated behind; margin of the valves crenulated.

In the British Museum. From Labrador; C. Goodsir.

TEREBRATULA ALGOENSIS. *Ter. testa suborbiculari, posticè subacuminata, anticè sublobata, albida; valvis radiatim striatis; carina dorsali conspicua, rotundata; foramine magno, incompleto; margine valvæ dorsali minutissimè crenulato.*

Shell suborbicular, slightly acuminated behind, rather lobed in front, whitish; valves radiately striated; mesial ridge distinct, roundish; perforation large, incomplete; margin of the dorsal valve very minutely crenulated.

A single valve of this specimen is in the British Museum, labelled "Algoa Bay, Bowerbank."

"Descriptions of new species of *Marginella*." By G. B. Sowerby, F.L.S.

MARGINELLA FUSCA. *Marg. testa elongata, subovali, posticè subangulata, anticè latè marginata, in medio paululum contracta, fuscâ, vel pallidè purpurea, fusco-trifasciata; spiræ breviuscula, apice obtuso; aperturæ angusta, columellæ rectiuscula, plicis quatuor, quarum duæ anticæ albæ, prominentibus, spiraliter elongatis; labio externo albo, intus in medio incurvo, extus fusco, latè reflexo.*

Differing from *M. nitida* in the colouring and the shortness of the spire, and in the outer lip being more broadly reflected.

In Mr. Cuming's collection. From the West Indies.

MARGINELLA CRASSILABRUM. *Marg. testá subovali, in medio subangulatā, pallidè griseo-fulvā; spirā brevi; anfractibus distinctis, ultimo dilatato, ad spiram elevato; columellā plicis quatuor, quarum duæ anticæ prominentes, spiraliter elongatis; labio externo, crasso, latè incurvo, angulato, extus varicoso, ad apicem tumidè elevato.*

This species is remarkable for the broad angular disc formed by the outer lip, which is much thickened at the back and raised so as nearly to cover the spire.

In Mr. Jackson's and Mr. Cuming's collections. From the West Indies.

MARGINELLA TÆNIATA. *Marg. testá ovali, cylindricā, pallidè fulvā, fasciis fuscis tribus cinctā; spirā brevi; aperturā elongatā, posticè subangustatā; columellā plicis quatuor, quarum duæ anticæ majores; labio externo lævi, latè reflexo.*

Differing from *M. avena* in having a shorter spire and the outer lip more broadly reflected.

In Mr. Cuming's collection. Locality unknown.

MARGINELLA ALBO-CINCTA. *Marg. testá subconicā, subangulatā, lævi; spirā productā; anfractibus angulatis, ultimo fascia alba prope angulum, et altero ad terminum anticū cincto inter fascias fusco maculato et punctis nigris picta; columellā quadriplicata.*

Provisionally described from a young specimen in Mr. Cuming's collection. When full-grown it would probably resemble *M. nubeculata* in form.

MARGINELLA PSEUDO-FABA. (*M. Faba*, Lam. Anim. s. vert., vii.)

Marg. testā angulatā, anticē attenuatā, subrecurvā, pallidè fulvā, griseo-nebulatā, punctorum irregularium seriebus 10 sparsim cinctā; spirā prominula; anfractibus angulatis, ad angulum validè crenulatis, crassis, paululum arcuatis, posticè angulatis, anticē emarginatis, attenuatis.

Much more angular than the true *M. Faba*, and has the anterior part of the body-whorl tapering and bent upwards.

In Mr. Cuming's collection. From the river Gambia, West Africa.

MARGINELLA FAUNA. *Marg. testā ovali, subcylindricā, pallidissimè carneā, spirā brevi; columellā obliquè quadriplicata; labio prope medium incurvo, extus subincrassato.*

Slightly resembling *M. pallida*, but more oval; the lower part of the aperture less open, and the outer lip thicker.

In Mr. Cuming's collection. From the isle of Curasso.

MARGINELLA MULTILINEATA. *Marg. testā ovali, straminea, lineis rubris numerosis cinctā; spirā penè celatā, apice fasciā rubra circulari cincto; aperturā anticē et posticē emarginata; columellā albā,*

in medio tumida, anticè callosa, plicis quatuor ad quinque inæqualibus; labio externo albo, intus crenulato, in medio subangulato, extus tenuiter reflexo.

On the whole resembling *tessellatus*, but it is much shorter, with the outer lip less varicose on the outside, and coloured by numerous red lines instead of the square patches.

In Mr. Cuming's collection. From Belize, bay of Honduras; Mr. Dyson.

MARGINELLA VARIA. *Marg. testā elongatā, lævi, anticè expansā, albā, vel fusca, vel fusco vel rubro trifasciatā, vel purpureo longitudinaliter et spiraliter interruptim fasciatā; spirā plus minusve productā; aperturā posticè angustā, anticè subexpansā; columella quadriplicatā; labio externo in medio incurvo, extus leviter varicoso.*

Differing from *M. lactea* of Kiener in being wider at the anterior termination, and in the outer lip not being so much elevated.

From the West Indies. Varieties are from Belize, bay of Honduras.

MARGINELLA SIMILIS. *Marg. testā ovali, straminea, griseo-nebulatā, lineis creberrimis interruptim cinctā; spirā penè celatā; aperturā angustā, anticè et posticè emarginatā; columellā spirā tumida, anticè varicosā, irregulariter septemplicatā; labio externo posticè spiram paululum superante, intus crenulato, extus nigro maculato vix marginato.*

The spire is less concealed, and the margin of reflected lip less distinct than in *M. interrupta*. The colouring is more mottled.

In Mr. Cuming's collection. From the Brazils.

October 27, 1846.

Richard C. Griffith, Esq., in the Chair. .

No business was transacted.